

Judson R. Roberts, DDS
Robin H. Berrin, DDS
Judith V. Lane, DDS
Robert W. Pretel, DDS, MSD

Capitol
PERIODONTAL
G R O U P
Periodontics & Implant Dentistry

P. Paul Towfighi, DDS, MS
P. Kevin Chen, DMD, MS
Rosemary Wu, DMD, MS

CARE OF YOUR MOUTH FOLLOWING PERIODONTAL SURGERY

Resting the day of your surgery and following these instructions will help ensure early, comfortable and complete healing.

1. DISCOMFORT – There may be some discomfort following your surgical procedure. To help control the discomfort, **please take your pain medications as soon as possible**. Eating prior to taking narcotic medications (such as Vicodin or Tylenol with codeine) is important to reduce the chance of nausea. **Please avoid taking aspirin or aspirin-containing compounds** and take your antibiotic on an empty stomach. If you are still in pain, you may alternate the narcotic pain medication every 2-3 hours with Ibuprofen (Motrin, Advil, etc.) if you are able to take it. This may reduce the amount of narcotic medication you'll need to take.

2. SWELLING – Swelling may accompany surgery. It may be partially controlled by applying ice packs as soon as possible and continuing the first day. The packs may be applied for 15 minutes on and 15 minutes off. After 24 hours, warm moist compresses should be applied to help reduce swelling. Ibuprofen may also reduce swelling faster.

3. BLEEDING – Slight oozing of blood may occur for the first 24 hours. If excessive or continuous bleeding occurs, discontinue any rinsing and avoid exertion. Prolonged bleeding can usually be controlled at home by using a moist gauze pad or a moist tea bag (Black Tea only), **applied with firm finger pressure** to the bleeding area for 20 minutes

or more. Pressure is key. **AVOID EXCESSIVELY HOT LIQUIDS THE FIRST SIX HOURS AFTER SURGERY**, as this can cause bleeding. Also, avoid spicy or acidic foods. **DO NOT DRINK THROUGH A STRAW. WHEN UNCONTROLLED BLEEDING PERSISTS, PLEASE CONTACT US OR GO TO THE NEAREST EMERGENCY ROOM.**

4. RINSING AND BRUSHING – After 24 hours you may rinse gently with warm salt water or mouthwash; however, people with high blood pressure should avoid salt. Please begin gentle brushing after 24 hours, using warm water.

5. SMOKING – Please do not smoke, or at least greatly restrict your smoking following surgery, to help improve your healing.

6. DIET – Diet should usually be limited to soft or semi-soft foods for the first few days. **It is important to maintain a good diet**. Nourishing and easy-to-eat foods include milkshakes, cottage cheese, yogurt, soup, ground meat, etc. Nutritional supplements such as Meritene, Nutriment, Ensure, etc. may be used for one or two meals.

7. DRESSING – A dressing or bandage may be used to cover the surgical area. It is of no concern if small pieces of the dressing break off after a few days, unless the area becomes uncomfortable.

**IF YOU HAVE ANY QUESTIONS, PLEASE DO NOT HESITATE TO CALL.
A DOCTOR IS ON CALL TO HELP YOU AT ANY TIME.
THE AFTER-HOURS EMERGENCY NUMBER IS (916) 971-3461.**

Sacramento Office
1810 Professional Drive
Sacramento, California 95825
(916) 971-3461
(916) 973-9830 Fax

Laguna Office
9309 Office Park Circle, Suite 110
Elk Grove, California 95758
(916) 684-3379
(916) 684-4106 Fax

Roseville Office
2428 Professional Drive, Suite 100
Roseville, California 95661
(916) 786-6585
(916) 786-7542 Fax

Folsom Office
2535 East Bidwell, Suite 150
Folsom, California 95630
(916) 984-1109
(916) 984-1764 Fax