

Judson R. Roberts, DDS
Robin H. Berrin, DDS
Judith V. Lane, DDS
Robert W. Pretel, DDS, MSD



P. Paul Towfighi, DDS, MS
P. Kevin Chen, DMD, MS
Rosemary Wu, DMD, MS

PREPARING YOURSELF FOR PERIODONTAL SURGERY

INTRAVENOUS SEDATION

Please do not eat or drink anything for six hours prior to surgery. However, ***please take any prescribed medications*** unless instructed otherwise. If you are scheduled for an afternoon intravenous sedation surgery, please get up early enough to eat breakfast at least six hours prior to surgery. **A responsible adult must take you home and remain with you following your surgery.**

LOCAL ANESTHESIA

No restrictions. You may eat and drink prior to surgery, and drive to and from the office. We encourage you to eat a light snack before your visit. Please do not skip meals.

IMPORTANT THINGS TO FOLLOW

- Any aspirin or anti-coagulant treatment will be addressed on an individual basis by your doctor prior to treatment.
- Please do not donate blood for 3-4 weeks prior to surgery.
- ***No alcohol*** consumption 24 hours prior to surgery.
- ***Take your normal PRESCRIPTION MEDICATION, unless instructed otherwise.***
- Notify staff of any medications that you take the day of your surgery.
- ***Wear a short-sleeve garment the day of your surgery.***
- Wear comfortable walking shoes. No high heels.
- If you are having intravenous sedation surgery, remove nail polish.
- Daily wear contact lenses should be removed prior to your surgery.
- Prepare ice packs and have soft nutritious foods available at home. Nutritious, easy to eat foods include: milkshakes, cottage cheese, yogurt, soup, ground meat, etc. Nutritional supplements such as Meritene, Nutriment or Ensure may also be consumed.

SPECIAL INSTRUCTIONS FOR DIABETICS

If using local anesthesia, eat and take medications as normal.

Insulin dependent diabetics undergoing IV sedation should take ½ their normal insulin dose if the surgery is in the morning. If the surgery is in the afternoon, then eat a normal breakfast and take your normal dose of insulin 6 hours before the surgery.

RECUPERATION TIME

The average recuperation period is 3-7 days, although some people require additional time. Other people do not require any time off from work. There is no mandatory length of bed rest required.

DATE OF SURGERY: _____

DATE OF SUTURE REMOVAL: _____

***Our staff and your doctor are available to answer any questions.
Please do not hesitate to call. The after-hours emergency number is (916) 971-3461***

Sacramento Office
1810 Professional Drive
Sacramento, California 95825
(916) 971-3461
(916) 973-9830 Fax

Laguna Office
9309 Office Park Circle, Suite 110
Elk Grove, California 95758
(916) 684-3379
(916) 684-4106 Fax

Roseville Office
2428 Professional Drive, Suite 100
Roseville, California 95661
(916) 786-6585
(916) 786-7542 Fax

Folsom Office
2535 East Bidwell, Suite 150
Folsom, California 95630
(916) 984-1109
(916) 984-1764 Fax