

Dental Practice of
Robin H. Berrin, DDS
Robert W. Pretel, DDS, MSD
P. Paul Towfighi, DDS, MS
P. Kevin Chen, DMD, MS



Rosemary Wu, DMD, MS
Nidhi Jain, BDS, DMD, MS
Aneel Nath, DDS
Khalid Rasheed, DDS

Halcion (Triazolam), or Valium (Diazepam) Instructions

If you have been prescribed Halcion or Valium for sedation, please follow the directions listed below:

- Take your medication as directed on the bottle.
- Please do not eat anything 4-6 hours prior to the procedure. You can drink clear fluids like water or apple juice.
- You must have a driver who stays with you from the time you take the sedation medication at home prior to your appointment, through a time 2-3 hours after you have returned home. **This means your driver must bring you to the appointment and come into the office with you.**
- **Your driver must stay at the office during your appointment and stay with you at least 2-3 hours after you've returned home.**
- **You may not drive yourself to and from your appointment.**
- **You may not drive for 24 hours after taking the medication.**

DO NOT use Halcion or Valium if:

- You have a severe mental disorder, angle-closure glaucoma, or severe liver disease.
- You are taking clozapine, delavirdine, efavirenz, HIV protease inhibitors (eg, ritonavir), itraconazole, ketoconazole, nefazodone, or sodium oxybate (GHB).

Precautions:

Halcion or Valium may interfere with certain medical conditions. Advise your doctor or pharmacist if the following medical conditions apply to you:

- Pregnant, planning to become pregnant, or are breast-feeding.
- History or current alcoholism or substance abuse/dependence, or if you drink alcohol.
- Depression, mental or mood problems, or have a history of suicidal thoughts or behavior.
- Myasthenia gravis (a condition of progress muscle paralysis).
- Liver or kidney problems, the blood disease porphyria, or glaucoma or you are at risk for glaucoma.
- Chronic bronchitis, chronic obstructive pulmonary disease (COPD), emphysema, difficulty breathing while asleep (sleep apnea), or other breathing problems.

Please Note: Excessive grapefruit or grapefruit juice consumption may prolong the sedative side effects of Halcion.

Sacramento Office

1810 Professional Drive, Suite B
Sacramento, California 95825
(916) 971-3461
(916) 973-9830 Fax
SacramentoAdmin@Capperio.com

Laguna Office

9309 Office Park Circle, Suite 110
Elk Grove, California 95758
(916) 684-3379
(916) 684-4106 Fax
LagunaAdmin@Capperio.com

Roseville Office

2428 Professional Drive, Suite 100
Roseville, California 95661
(916) 786-6585
(916) 786-7542 Fax
RosevilleAdmin@Capperio.com

Folsom Office

2535 East Bidwell, Suite 150
Folsom, California 95630
(916) 984-1109
(916) 984-1764 Fax
FolsomAdmin@Capperio.com